

AN EFFICIENT METHOD OF FINDING A PUISEUX EXPANSION OF A PARAMETRIC SINGULARITY

MACIEJ BORODZIK

ABSTRACT. We present a new and very efficient method of determining a topological type of a cuspidal plane curve singularity parametrised locally by $(x(t), y(t))$. The approach relies on defining inductively a sequence of functions P_k which are polynomials in x , y and derivatives of x and y .

Let us be given a germ of a parametric curve in $\mathbb{C}^2$ given by

$$(1) \quad \begin{aligned} x(t) &= a_0 t^p + a_1 t^{p+1} + \dots, \\ y(t) &= b_0 t^q + b_1 t^{q+1} + \dots \end{aligned}$$

Here $a_0 b_0 \neq 0$, $q \geq p > 1$ and both series on the right hand side are assumed to be convergent in some neighbourhood of $t = 0$. In order to determine the topological type of the singularity (1) we need (see [BK, EN]) to find its characteristic sequence $(p; q_0, q_1, \dots, q_n)$. To do this we write the Puiseux expansion of (1) in the form

$$(2) \quad y = c_q x^{q/p} + c_{q+1} x^{(q+1)/p} + \dots$$

Then we define $q_0 = \min\{s \geq 0: c_s \neq 0 \wedge p \nmid s\}$, and inductively $p_0 = p$, $p_k = \gcd(p_{k-1}, q_{k-1})$. If $p_k \neq 1$ we define $q_k = \min\{s \geq 0: c_s \neq 0 \wedge p_k \nmid s\}$; if $p_k = 1$ we stop the procedure. As $p_k > p_{k+1}$ this procedure will definitely stop after a finite number of steps. So let n be such that $p_{n+1} = 1$, then $(p; q_0, \dots, q_n)$ is the characteristic sequence.

Therefore, in order to know the topological type of singularity we need to know the coefficients c_k , or, at any rate, to know which one are zero and which are not. Now passing from (1) to (2) is, in general, a very involved procedure, which requires finding an explicit formula for $t = d_1 x^{1/p} + d_2 x^{2/p} + \dots$. Moreover, the Puiseux coefficients c_k are not very well defined, namely they are functions of some fractional powers of a_0 . In particular if we study a deformation, in other words if we allow the coefficients of (1) to vary with some parameter s , and $a_0(s) \rightarrow 0$ as $s \rightarrow s_0$, then $c_k(s)$ can easily get out of hand. Some of them can diverge to infinity, some of them can converge

Date: 8 Dec 2009.

2000 Mathematics Subject Classification. Primary 14H20; Secondary 14Q05.

Key words and phrases. plane curve singularity, Puiseux expansion, characteristic sequence.

Supported by Polish MNiSz Grant N N201 397937. The author is also supported by Foundation for Polish Science (FNP).

but lose any topological meaning. We want to propose a remedy to both problems: we shall define some polynomials in y , x and their derivatives over t , which carry the same data about the singularity as Puiseux expansion and such that one can recover the Puiseux expansion from them in a very easy way.

As a matter of fact, the definition of polynomials P_k appears first in [Bo], but only after writing this article the author realised that the polynomials P_k may be of wider use and allow fast and efficient computation of a characteristic sequence.

We begin with putting

$$P_0(t) = y \text{ and } r_0 = \text{ord}_{t=0} P_0(t) = q$$

Like in [BZ1, Proof of Lemma 3.2], let us divide (2) by $x^{q/p}$ and differentiate both sides with respect to t (it is convenient to write here $\dot{x} = \frac{dx}{dt}$ and so on). We get

$$\frac{\dot{y}x - \frac{q}{p}\dot{x}y}{x^{q/p+1}} = \frac{1}{p}c_{q+1}\dot{x}x^{1/p-1} + \frac{2}{p}c_{q+2}\dot{x}x^{2/p-1} + \dots$$

So if

$$(3) \quad P_1 = \dot{y}x - \frac{q}{p}\dot{x}y$$

we will have

$$P_1 = \frac{1}{p}c_{q+1}x^{(q+1)/p}\dot{x} + \frac{2}{p}c_{q+2}x^{(q+2)/p}\dot{x} + \dots$$

Let

$$(4) \quad r_1 = \text{ord}_{t=0} P_1 - (p-1).$$

Then we have an obvious lemma.

Lemma 1. *The coefficients $c_{r_0+1}, \dots, c_{r_1-1}$ all vanish. Moreover the first not vanishing coefficient of P_1 , i.e. the one at $t^{r_1+(p-1)}$, is equal to $(r_1 - r_0)c_{r_1} \cdot a_0^{r_1/p+1}$.*

Now we define inductively the polynomials P_k and the numbers r_k . The induction procedure is as follows. Assume that we have

$$(5) \quad P_k = \frac{r_k - r_{k-1}}{p} \cdot \frac{r_k - r_{k-2}}{p} \cdot \dots \cdot \frac{r_k - r_0}{p} c_{r_k} \dot{x}^{2k-1} x^{r_k/p} + \dots$$

with

$$(6) \quad r_k = \text{ord}_{t=0} P_k - (2k-1)(p-1).$$

In (5) dots denote terms like $(s - r_{k-1}) \dots (s - r_0) p^{-k} c_s \dot{x}^{2k-1} x^{s/p}$, which are too long to be written in a nice looking formula.

Now we divide both sides of (5) by $\dot{x}^{2k-1} x^{r_k/p}$, differentiate both sides with respect to t and then multiply them again by $\dot{x}^{2k+1} x^{r_{k+1}/p+1}$. We get

$$P_{k+1} = \frac{r_{k+1} - r_k}{p} \frac{r_{k+1} - r_{k-1}}{p} \cdot \dots \cdot \frac{r_{k+1} - r_0}{p} c_{r_{k+1}} \dot{x}^{2k+1} x^{r_{k+1}/p} + \dots,$$

where

$$(7) \quad r_{k+1} = \text{ord}_{t=0} P_{k+1} - (2k+1)(p-1)$$

and

$$(8) \quad P_{k+1} = x\dot{x}\frac{d}{dt}P_k - \left(\frac{r_k}{p}\dot{x}^2 + (2k-1)\ddot{x}x\right)P_k.$$

Similarly we have a lemma the proof of which is an easy computation.

Lemma 2. *The Puiseux coefficients $c_{r_{k-1}+1}, \dots, c_{r_k-1}$ all vanish. The coefficient at $t^{r_k+(2k-1)(p-1)}$ of P_k , i.e. the first non-vanishing one, is equal to*

$$(r_k - r_{k-1})(r_k - r_{k-2}) \dots (r_k - r_0)p^{k-1}c_{r_k}a_0^{r_k/p+(2k-1)}.$$

Thus, using the above method we can easily find the Puiseux expansion. The computations become simpler if there are many Puiseux terms that vanish. The algorithm, in a form easy to use, can be presented as follows.

Algorithm. Given a parametric presentation as in (1) of a cuspidal singularity with $a_0b_0 \neq 0$.

- Step 1. Set $p_0 = p$ and $r_0 = q$. Let $p_1 = \gcd(p_0, q_0)$. If $p_1 = 1$ then the singularity is quasi-homogeneous and we stop.
- Step 2. Compute P_1 as in (3) and r_1 from (4). Let $p_2 = \gcd(p_1, r_1)$.
- Step 3. If $p_2 = 1$ then we stop. Otherwise put $k = 1$.
- Step 4. Compute P_{k+1} and r_{k+1} from (8) and (7) respectively.
- Step 5. Let $p_{k+1} = \gcd(p_k, r_k)$ if $p_{k+1} = 1$ we stop, otherwise we increase k and go to Step 4.

As an output we get a sequence of positive integers $(r_0, r_1, r_2, \dots, r_n)$, such that the Puiseux expansion is

$$y = c_{r_0}x^{r_0/p} + c_{r_1}x^{r_1/p} + \dots + c_{r_k}x^{r_k/p} + \text{inessential Puiseux terms}.$$

Here $c_{r_0}, c_{r_1}, \dots, c_{r_k}$ are all non-zero. The procedure will eventually stop unless the parametrisation (1) is not one to one.

Of course, we can recover not only the characteristic sequence, but also the exact values of non-vanishing Puiseux coefficients. It is enough to look at the first non-vanishing coefficients of functions P_{k+1} and use second part of Lemma 2.

Example. Consider a singularity parametrised by

$$\begin{aligned} x(t) &= t^{12} + t^{13} + \frac{37}{28}t^{14} \\ y(t) &= t^{18} + \frac{3}{2}t^{19} + \frac{33}{14}t^{20} + \frac{13}{14}t^{21} + \frac{675}{1568}t^{22} - \frac{675}{3136}t^{23}. \end{aligned}$$

We write

$$y = x^{18/12} + c_{19}x^{19/12} + \dots$$

But it is easy to compute that

$$P_1 = -\frac{2025}{10976}t^{35} - \frac{24975}{43904}t^{36}.$$

So $r_1 = 35 - 11 = 24$. Thus $c_{19} = c_{20} = c_{21} = c_{22} = c_{23} = 0$ and $c_{24} \neq 0$. Then

$$P_2 = \frac{2500875}{76832}t^{59} + \dots$$

so $r_2 = 59 - 3 \cdot 11 = 26$, so $c_{25} = 0$ and $c_{26} \neq 0$. Then a simple computation shows that $P_3 \sim t^{82}$, so $c_{27} \neq 0$ and the Puiseux expansion is $y = x^{3/2} + c_{24}x^2 + c_{26}x^{13/6} + c_{27}x^{27/12} + \dots$, so the characteristic sequence is $(12; 18, 26, 27)$ (c_{24} is inessential).

We can use the polynomials P_k in studying deformations. In fact, they behave very well if $x(t)$ and $y(t)$ vary smoothly. This approach is pursued in [Bo] with quite a success in some partial cases.

Acknowledgement. *The author wishes to express his thanks for H. Żołądek, A. Płoski and P. Cassou-Noguès for fruitful discussions and to Christian Gorzel for his many comments to the first version of the paper.*

REFERENCES

- [BK] E. Brieskorn, H. Knörrer, *Plane Algebraic Curves*, Birkhäuser, Basel-Boston-Stuttgart, 1986.
- [Bo] M. Borodzik *Puiseux coefficients and parametric deformations of plane curve singularities*, preprint, [arxiv:0909.4469](https://arxiv.org/abs/0909.4469).
- [BZ1] M. Borodzik, H. Żołądek *Geometry of Puiseux expansions*, Ann. Polon. Math. **93** (2008), no. 3, 263–280.
- [EN] D. Eisenbud and W. Neumann, *Three-dimensional link theory and invariants of plane curve singularities*, Annals Math. Studies **110**, Princeton Un-ty Press, Princeton, 1985.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097
WARSAW, POLAND
E-mail address: mcboro@mimuw.edu.pl